

#### STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers.

#### All other non-refillable containers

Triple rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product. **DO NOT** re-use empty container for any other purpose.

#### Plastic Bags and cardboard containers

Shake empty bag into spray tank. Single rinse before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. Break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty bags or product.

Dispose of spent dip in an authorised dip disposal facility. If an authorised dip disposal facility is not available, the spent dip should be evenly spread over flat land not exceeding 20,000L/ha. The disposal site must be dedicated, limed and adequately bunded (soil at least 15cm high). **DO NOT** dispose unwanted spent dip in the same place for at least 420 days, as repeated depositions in one location may, over time, create a contaminated site. Unused or spent dips should be disposed of carefully to avoid contamination of wetlands or watercourses.

#### SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing product for use, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. When using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). Wash hands after use. After each day's use, wash gloves, and contaminated clothing.

# DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

IMTRADE

**NOSCLEX<sup>®</sup>**  
800  
WG FUNGICIDE

ACTIVE CONSTITUENT: 800 g/kg PROCYMIDONE

**GROUP 2 FUNGICIDE**

For the control of certain fungal diseases on various crops  
as specified in the Directions for Use table

**IMPORTANT: Read the attached Leaflet before use.**

FORM  
**WG**

CONTENTS  
**1kg-10kg**



Batch No.  
DOM:

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre  
(Phone Australia 13 11 26; New Zealand 0800 764 766).

Refer to attached Leaflet for additional GHS  
Hazard & Precautionary Statements

#### SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from your supplier or Imtrade Australia Pty Ltd website at [www.imtrade.com.au](http://www.imtrade.com.au)

#### CONDITION OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

NOSCLEX<sup>®</sup> is a Registered Trademark of Imtrade Australia Pty Ltd

In a Transport Emergency Dial 000 Police or Fire Brigade

APVMA Approval No: 70284/RV22

**IMTRADE**  
CropScience

17 Ocean St, Kwinana Beach WA 6167  
08 9419 0333 | [www.imtrade.com.au](http://www.imtrade.com.au)

**DANGEROUS POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY INSTRUCTIONS BEFORE OPENING OR USING



# **IMTRADE NOSCLEX® 800 WG FUNGICIDE**

**ACTIVE CONSTITUENT: 800 g/kg PROCYMIDONE**

**GROUP**

**2**

**FUNGICIDE**

For the control of a certain fungal diseases on various crops  
as specified in the Directions for Use table

**IMPORTANT: READ THIS LEAFLET THOROUGHLY  
BEFORE OPENING AND USING THIS PRODUCT**

**APVMA Approval No: 70284/RV22**

Imtrade Australia Pty Ltd  
ABN 13 090 151 134  
17 Ocean Street, Kwinana WA 6167  
Tel: 1800 171 799 Fax: 1800 171 788  
Web: [www.imtrade.com.au](http://www.imtrade.com.au)

## RESTRAINTS:

**DO NOT** apply to Couch Grass greens in the period April-September, or when Couch Grass is near dormancy.

**DO NOT** use Imtrade NOSCLEX® 800 WG Fungicide in the home garden.

## SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at [www.apvma.gov.au/spraydrift](http://www.apvma.gov.au/spraydrift)

- **DO NOT** allow **bystanders** to come into contact with the spray cloud.
- **DO NOT** apply in a manner that may cause an unacceptable impact to **native vegetation, agricultural crops, landscaped gardens** and **aquaculture production**, or cause contamination of plant or livestock commodities, outside the application site from **spray drift**. The **buffer zones** in the relevant buffer zone tables below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.
- **DO NOT** apply unless **wind speed** is between 3 and 20 kilometres per hour at the **application site** during the time of application.
- **DO NOT** apply if there are **hazardous surface temperature inversion** conditions present at the **application site** during the time of application. **Surface temperature inversion conditions** exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

## BOOM SPRAYERS

- **DO NOT** apply by a boom sprayer unless the following requirements are met:
  - o Spray droplets are not smaller than a **MEDIUM spray droplet size category**.
  - o Minimum distances between the **application site** and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer Zones for Boom Sprayers') are observed.

### Buffer zones for Boom sprayers

Application Rate	Boom height above the target Canopy	Mandatory buffer zones				
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
Up to 6kg/ha	0.5m or lower	0m	10m	0m	0m	350m
	1.0m or lower	0m	30m	0m	0m	375m
Up to 4kg/ha	0.5m or lower	0m	0m	0m	0m	200m
	1.0m or lower	0m	25m	0m	0m	350m
Up to 3.5kg/ha	0.5m or lower	0m	0m	0m	0m	160m
	1.0m or lower	0m	20m	0m	0m	350m
Up to 1.25kg/ha	0.5m or lower	0m	0m	0m	0m	30m
	1.0m or lower	0m	10m	0m	0m	140m
Up to 600g/ha	0.5m or lower	0m	0m	0m	0m	10m
	1.0m or lower	0m	0m	0m	0m	60m
Up to 300g/ha	0.5m or lower	0m	0m	0m	0m	0m
	1.0m or lower	0m	0m	0m	0m	30m

## VERTICAL SPRAYERS

**DO NOT** apply by a vertical sprayer unless the following requirements are met:

- o Spray is not directed above the target canopy.
- o The outside of the sprayer is turned off when turning at the end of rows and when spraying the outer row on each side of the application site.
- o For dilute water rates up to the maximum listed for each type of canopy specified, minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for vertical sprayers') are observed.

### Buffer zones for Vertical sprayers

Type of target canopy and dilute water rate	Mandatory buffer zones				
	Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
2m tall and shorter, maximum dilute water rate of 1000L/ha	0m	0m	0m	0m	0m
Taller than 2m (not fully-foliated), maximum dilute water rate of 1500L/ha	0m	0m	0m	0m	20m
Taller than 2m (fully-foliated), maximum dilute water rate of 1500L/ha	0m	0m	0m	0m	10m

## AIRCRAFT

**DO NOT** apply by aircraft unless the following requirements are met:

- o Spray droplets not smaller than a MEDIUM **spray droplet size category** for Lentil application.
- o Spray droplets not smaller than a COARSE **spray droplet size category** for Canola application.
- o For maximum release height above the target canopy of 3 m or 25% of wingspan or 25% of rotor diameter, whichever is the greatest, minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for aircraft') are observed.

### Buffer zones for Aircraft (MEDIUM spray droplet size)

Application Rate	Boom height above the target Canopy	Mandatory buffer zones				
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
Up to 300g/ha	Fixed-wing	0m	0m	0m	0m	220m
	Helicopter	0m	10m	0m	0m	140m

### Buffer zones for Aircraft (COARSE spray droplet size)

Application Rate	Boom height above the target Canopy	Mandatory buffer zones				
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
Up to 300g/ha	Fixed-wing	0m	0m	0m	0m	220m
	Helicopter	0m	10m	0m	0m	140m

## DIRECTIONS FOR USE

Crop	Disease Controlled	States	Application Rate	WHP	Critical Comments
Canola	Sclerotinia Rot ( <i>Sclerotinia sclerotiorum</i> )	All States	600g/ha <b>Ground Application</b> in 100L/ha water plus Agral® at 20mL/100L water <b>Aerial Application</b> in minimum 40L/ha water plus Agral® at 20mL/100L water	Nil (H) 9 weeks (G)	<p>Spraying should occur before petals begin to drop and preferably prior to a rainfall event during the early-mid flowering stage of crop growth.</p> <p>Infection of Canola stems and branches occurs when infected petals fall and lodge in the lower canopy of the plant, particularly during wet or humid conditions.</p> <p>The objective of applying Imtrade NOSCLEX® 800 WG Fungicide is to treat as many petals as possible prior to petal drop and before pods set.</p> <p>Application should, therefore take place by 30% bloom (i.e. 30% of flowers open on the main stem), at which stage the maximum number of flowers are open at one time and little petal fall has occurred.</p> <p>Application should not be made after mid-flowering.</p> <p>For aerial application to a Canola crop that is directly adjacent to a downwind paddock where grazing stock may be present, use coarse nozzle settings and observe a 20m buffer to that paddock in order to minimise drift.</p>

Crop	Disease Controlled	States	Application Rate	WHP	Critical Comments
Lentils	Grey Mould ( <i>Botrytis cinerea</i> and <i>Botrytis fabae</i> )	All States	300g/ha Apply in a minimum of 100L/ha water for <b>ground application</b> or 45L/ha for <b>aerial application</b>	21 days (H) 21 days (G)	<p>Monitoring of crops for disease should commence at 6-8 weeks after crop emergence. Early application of fungicide is critical in restricting the development and spread of Grey Mould.</p> <p>The first application of Imtrade NOSCLEX® 800 WG Fungicide is recommended immediately <b>prior to canopy closure</b> to ensure good spray penetration into the crop. Subsequent monitoring of crop and environmental conditions will help determine timing of later applications.</p> <p>Other critical growth stages for disease control are:</p> <ul style="list-style-type: none"> <li>▶ mid-flowering/early pod fill</li> <li>▶ end of flowering/late pod fill</li> </ul> <p>Later fungicide applications may be required if conditions are conducive to disease development.</p> <p>Apply no more than two consecutive sprays of Imtrade NOSCLEX® 800 WG Fungicide. Alternate with fungicides with different Modes of Action. Imtrade NOSCLEX® 800 WG Fungicide will not provide effective control of Ascochyta Blight (<i>Ascochyta lentis</i>).</p> <p>For aerial application use medium to coarse nozzle settings in order to minimize drift.</p>

Crop	Disease Controlled	States	Application Rate	WHP	Critical Comments
Grapes (Wine Grapes only)	Grey Mould ( <i>Botrytis cinerea</i> )	All States & NT	<b>Dilute Spraying</b> 45g/100L <b>Concentrate Spraying</b> Refer to the <b>Mixing/ Application</b> section	9 days	<b>DO NOT</b> use on table grapes or grapes used for the production of dried fruit. Use on wine grapes only. Apply at the following growth stages: <ul style="list-style-type: none"> <li>➤ 80% cap fall,</li> <li>➤ just prior to bunch closure,</li> <li>➤ at veraison (when sugar content rises),</li> <li>➤ and 2-3 weeks pre-harvest.</li> </ul> To ensure complete bunch wetting add Agral® at 10-20 mL/100L. Apply by dilute or concentrate spraying equipment. Apply the same total amount of Imtrade NOSCLEX® 800 WG Fungicide to the target crop whether applying Imtrade NOSCLEX® 800 WG Fungicide by dilute or concentrate spraying methods. <b>DO NOT</b> use at concentrations greater than 95gm/100L of water.
Stone Fruit	Blossom Blight ( <i>Monilinia laxa</i> )	VIC, NSW, SA, TAS & QLD only	<b>Dilute Spraying</b> 30 to 45g/100L <b>Concentrate Spraying</b> Refer to the <b>Mixing/ Application</b> section	9 days	Apply at 10% blossom, full bloom, late petal and shuck fall. <b>DO NOT</b> apply after shuck fall. If weather conditions particularly favour Blossom Blight use higher rate. <b>NSW, SA, QLD &amp; TAS only:</b> Where <i>Monilinia laxa</i> is known to occur apply an additional early spray at pink bud. Apply by dilute or concentrate spraying equipment. Apply the same total amount of Imtrade NOSCLEX® 800 WG Fungicide to the target crop whether applying Imtrade NOSCLEX® 800 WG Fungicide by dilute or concentrate spraying methods. <b>DO NOT</b> use at concentrations greater than 95gm/100L of water.

Crop	Disease Controlled	States	Application Rate	WHP	Critical Comments
Onions	White Rot ( <i>Sclerotium cepivorum</i> )	VIC, QLD, NSW, SA & NT only	1.25kg/ha in a minimum 250L of water	4 weeks	<p><b>SOIL SPRAY:</b></p> <p>(a) Apply to soil surface immediately after sowing and repeat application at 10 weeks after sowing.</p> <p>(b) A further soil spray of 1.25kg/ha may be necessary if frequent or extended periods of cool moist conditions occur later in the season.</p> <p><b>NOTE:</b></p> <p>1. <b>DO NOT</b> spray directly over exposed seed in furrows before covering with soil.</p> <p>2. <b>WARNING:</b> Soil persistence Imtrade NOSCLEX® 800 WG Fungicide can be reduced under alkaline soil conditions.</p>
		VIC, QLD, NSW & SA only	600g/100L of water	4 weeks	<p><b>TRANSPLANT DIP:</b></p> <p>(a) Dip seedlings for up to 4 hours in fungicide suspension before transplanting.</p> <p>(b) A supplementary soil spray of 1.25kg/ha may be necessary if frequent or extended periods of cool moist conditions occur later in the season.</p>
Potato	Sclerotinia ( <i>Sclerotinia minor</i> )	All States & NT	300g to 600g/ha	21 days	<p>Apply first spray just before hilling up. Apply a second spray just after hilling. Direct these sprays towards the stem bases and soil surface. Apply in sufficient water to ensure thorough coverage. Use higher rate in situations where high disease levels are expected.</p> <p>Supplementary applications of 600g/ha to foliage at 14-21 day intervals may be necessary if conditions favour further development of disease.</p>
	Target Spot ( <i>Alternaria solani</i> )		300g/ha	21 days	<p>Apply in a program of sprays at 10 day intervals, beginning when warm weather conditions favour the disease and plants are 150 to 190mm high. Apply in sufficient water to ensure thorough coverage.</p>



Crop	Disease Controlled	States	Application Rate	WHP	Critical Comments
Ornamentals	Sclerotinia Rot ( <i>Sclerotinia sclerotiorum</i> )	NSW only	45 or 60g/100L of water	–	Apply to run-off. Use the higher rate when disease is severe. <b>DO NOT</b> apply to open African Violet flowers.
Turfgrass	Dollar Spot ( <i>Sclerotinia homeocarpa</i> )	VIC, QLD, ACT, NSW, SA & WA only	40 – 60g per 100m <sup>2</sup>	<b>DO NOT</b> graze treated turf or lawn; or feed turf or lawn clippings from any treated area to poultry or livestock	Apply in 5-10L water per 100m <sup>2</sup> . Use the higher rate where conditions conducive to severe disease occur. Apply at first sign of disease and repeat applications at intervals of 3-4 weeks. <b>CAUTION: Note phytotoxicity warning in General Instructions.</b>
	Black Helminthosporium ( <i>Drechslera</i> sp. <i>Bipolaris</i> sp. <i>Exserohilum</i> sp.)	VIC, QLD, ACT, NSW, SA & WA only	40 – 60g per 100m <sup>2</sup>		Apply in 5-10L water per 100m <sup>2</sup> . Apply at first sign of disease. A second application may be required after 2 to 4 weeks. <b>NOTE:</b> Imtrade NOSCLEX® 800 WG Fungicide spray programme for Spring Dead Spot will give preventative control of Black Helminthosporium until April. <b>CAUTION: Note phytotoxicity warning in General Instructions.</b>
	Spring Dead Spot ( <i>Leptosphaeria namarī</i> )				Apply in 5-10L water per 100m <sup>2</sup> . Apply Imtrade NOSCLEX® 800 WG Fungicide as the first two sprays of a monthly programme of four sprays beginning in February. Switch to an alternative fungicide such as TMTD® for the April and May applications. <b>CAUTION: DO NOT</b> apply to hybrid couch varieties from April through to September. <b>Note phytotoxicity warning in General Instructions.</b>

(H): Harvest (G): Grazing or cutting for stock feed

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

## **WITHHOLDING PERIOD**

**ONIONS: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.**

**STONEFRUIT (BLOSSOM BLIGHT CONTROL), WINE-GRAPES: DO NOT HARVEST FOR 9 DAYS AFTER APPLICATION.**

**POTATOES: DO NOT HARVEST FOR 9 DAYS AFTER APPLICATION.**

### **CANOLA:**

**HARVEST – NOT REQUIRED WHEN USED AS DIRECTED.**

**GRAZING – DO NOT GRAZE OR CUT FOR STOCK FEED FOR 9 WEEKS AFTER APPLICATION.**

### **LENTILS:**

**HARVEST – DO NOT HARVEST FOR 21 DAYS AFTER LAST APPLICATION.**

**GRAZING – DO NOT GRAZE OR CUT FOR STOCK FEED FOR 21 DAYS AFTER LAST APPLICATION.**

**TURFGRASS: DO NOT GRAZE TREATED TURF OR LAWN; OR FEED TURF OR LAWN CLIPPINGS FROM ANY TREATED AREA TO POULTRY OR LIVESTOCK.**

## **EXPORT OF TREATED LENTILS**

Growers should note that suitable MRLs or import tolerances may not be established in all markets for Lentils treated with Procymidone. If you are growing produce for export, please check with your exporter or Imtrade CropScience for the latest information on MRLs and import tolerances before using Imtrade NOSCLEX® 800 WG Fungicide.

## **GENERAL INSTRUCTIONS**

### **PHYTOTOXICITY**

Imtrade NOSCLEX® 800 WG Fungicide may cause discolouration and reduced growth of Colonial Bentgrass cv. Browntop. Imtrade NOSCLEX® 800 WG Fungicide is phytotoxic to couch grasses when they are approaching dormancy or are in a semi-dormant state and especially under cold very wet soil conditions. The phytotoxicity may result in scorching, reduced growth and loss of turf through wear. Avoid using Imtrade NOSCLEX® 800 WG Fungicide on couch greens within the period April to September. When used as directed, Imtrade NOSCLEX® 800 WG Fungicide may cause slight and temporary discolouration of Hybrid Couch cv. Tifgreen and other Tif-related varieties.

### **Mixing**

Imtrade NOSCLEX® 800 WG Fungicide is suitable for application through conventional spray equipment calibrated to ensure thorough crop coverage. Add the required amount of Imtrade NOSCLEX® 800 WG Fungicide to the partly filled tank with the agitator running and complete filling the tank with water. **DO NOT** mix with alkaline water. Continue thorough agitation during spraying and after a stoppage. **DO NOT** let prepared spray solution sit in spray tank overnight.

### **Dilute Spraying**

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the required amount of Imtrade NOSCLEX® 800 WG Fungicide specified in the Directions for Use table for each 100L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

### **Concentrate Spraying**

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

#### **EXAMPLE ONLY:**

1. Dilute spray volume as determined above: **for example** 1500L/ha.
2. Your chosen concentrate spray volume: **for example** 500L/ha.
3. The concentration factor in this example is: 3X (that is, 1500L divided by 500L = 3)
4. If the dilute rate is 6g/100L, then the concentrate rate becomes 3 x 6g, that is, 18g/100L of concentrate spray.
  - The chosen spray volume, amount of Imtrade NOSCLEX® 800 WG Fungicide per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows.
  - **DO NOT** use a concentrate rate higher than that specified in the 'Critical Comments'.
  - For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow Industry Best Practices.

## Tank Mixtures

Read and follow all label directions including application rates and safety directions for the tank mix products.

## Wetting Agent

Add a non-ionic surfactant at the rate directed on the product label.

## Compatibility

Imtrade NOSCLEX® 800 WG Fungicide is compatible with the following products: Agral®, Ambush®, Imtrade Pirimicarb 500 WG Aphicide, Omite®.

Provided the spray tank has an adequate agitation system and the mixture is not allowed to stand without agitation, these mixtures are acceptable. Some settling will occur with mixtures of Omite® and Ambush®.

## FUNGICIDE RESISTANT WARNING

GROUP	<b>2</b>	FUNGICIDE
-------	----------	-----------

Imtrade NOSCLEX® 800 WG Fungicide is a member of the dicarboximide group of fungicides. For fungicide resistance management Imtrade NOSCLEX® 800 WG Fungicide is a Group 2 Fungicide. Some naturally occurring individual fungi resistant to Imtrade NOSCLEX® 800 WG Fungicide or any other Group 2 Fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Imtrade NOSCLEX® 800 WG Fungicide or other Group 2 Fungicides. Since the occurrence of resistant fungi is difficult to detect prior to use, Imtrade CropScience accepts no liability for any losses that may result from the failure of Imtrade NOSCLEX® 800 WG Fungicide to control resistant fungi.

## PRECAUTIONS

### Re-entry

**DO NOT** enter treated areas until spray has dried. If prior entry is necessary, wear cotton overalls buttoned to the neck and wrist, elbow-length chemical resistant gloves and goggles.

## PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

### Toxic to aquatic life.

**DO NOT** contaminate wetlands or watercourses with this product or used containers.

## **STORAGE AND DISPOSAL**

**The method of disposal of the container depends on the container type. Read the Storage and Disposal instructions on the label that is attached to the container.**

Keep out of reach of children. Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers.

### ***Containers 1kg***

Rinse containers before disposal. Add rinsings to the spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product.

### ***All other non-refillable containers***

Triple rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product. **DO NOT** re-use empty container for any other purpose.

### ***Plastic Bags and cardboard containers***

Shake empty bag into spray tank. Single rinse before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. Break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty bags or product.

Dispose of spent dip in an authorised dip disposal facility. If an authorised dip disposal facility is not available, the spent dip should be evenly spread over flat land not exceeding 20,000L/ha. The disposal site must be dedicated, limed and adequately bunded (soil at least 15cm high). **DO NOT** dispose unwanted spent dip in the same place for at least 420 days, as repeated depositions in one location may, over time, create a contaminated site. Unused or spent dips should be disposed of carefully to avoid contamination of wetlands or watercourses.

## SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing product for use, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. When using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). Wash hands after use. After each day's use, wash gloves, and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (*Phone Australia 13 11 26; New Zealand 0800 764 766*).

### **Additional GHS Hazard & Precautionary Statements**

- Harmful if swallowed; •Harmful if inhaled; •May damage fertility or the unborn child;
- Avoid breathing dusts; •Wash contacted areas thoroughly after handling; •**DO NOT** eat, drink or smoke when using this product; •Use only outdoors or in a well-ventilated area; •Wear protective gloves, protective clothing and eye or face protection; •**IF SWALLOWED:** Call a POISON CENTRE or doctor if you feel unwell; •**IF SWALLOWED:** Rinse mouth. **DO NOT** induce vomiting; •**IF ON SKIN:** Wash with plenty of soap and water; •**IF INHALED:** Remove victim to fresh air and keep at rest in a position comfortable for breathing; •If exposed or concerned: Get medical advice; •If skin irritation occurs: Get medical advice; •If eye irritation persists: Get medical advice;
- Collect spillage; •In case of fire, use carbon dioxide, dry chemical, foam, water fog.

## SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from your supplier or Imtrade CropScience website at [www.imtrade.com.au](http://www.imtrade.com.au)

## CONDITION OF SALE

Imtrade CropScience shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade CropScience has any authority to add to or alter these conditions.

**In a Transport Emergency  
Dial 000  
Police or Fire Brigade**

**APVMA Approval No: 70284/RV22**