

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area out of direct sunlight. Shake empty bag into spray tank. Single rinse plastic bags before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. Puncture or shred and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. **DO NOT** inhale dust or spray mist. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), a washable hat and elbow-length PVC gloves. If using a hand directed sprayer, wear in addition: waterproof trousers and boots. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26; New Zealand 0800 764 766).

Refer to attached Leaflet for additional GHS Hazard & Precautionary Statements

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or from Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

IMTRADE

ATRAZINE 900 WG

HERBICIDE

ACTIVE CONSTITUENT: 900g/kg ATRAZINE

GROUP **5** HERBICIDE

For the control of weeds and grasses in Sorghum, Maize, Sugar Cane, TT-Canola, Lucerne, Oil Tea Tree and for fallow area maintenance and other situations as specified in the Directions for Use table

IMPORTANT: Read the attached Leaflet before use

UN 3077	ENVIRONMENTALLY HAZARDOUS SUBSTANCES, SOLID, N.O.S. (contains ATRAZINE)
In a Transport Emergency Dial 000 Police or Fire Brigade	
PG III	HAZCHEM 2Z

APVMA Approval No: 63503/122726



GROUP
5

FORM
WG

CONTENTS
1kg-50kg



Batch No.
DOM:

IMTRADE
CropScience

17 Ocean St, Kwinana Beach WA 6167
08 9419 0333 | www.imtrade.com.au

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



IMTRADE ATRAZINE 900 WG HERBICIDE

ACTIVE CONSTITUENT: 900 g/kg ATRAZINE

GROUP	5	HERBICIDE
--------------	----------	------------------

For the control of weeds and grasses in Sorghum, Maize,
Sugar Cane, TT-Canola, Lucerne, Oil Tea Tree
and for fallow area maintenance and other situations
as per the Directions for Use Table

**IMPORTANT: READ THIS LEAFLET THOROUGHLY
BEFORE OPENING OR USING THIS PRODUCT**

APVMA APPROVAL No: 63503/122726

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana WA 6167
Tel: 1800 171 799 Fax: 1800 171 788
Web: www.imtrade.com.au

RESTRAINTS

DO NOT use as a pre-emergent spray on light sandy soils.

DO NOT use on weeds over 4cm tall.

DO NOT apply to waterlogged soil.

DO NOT apply if heavy rains or storms that are likely to cause surface runoff are forecast within two days of application.

DO NOT irrigate to the point of runoff for at least two days after application.

DO NOT use as a pre-emergence application in Sorghum, Broom Millet or Saccaline during the wet season in the northern irrigation areas of Western Australia.

DO NOT apply product to any drainage line. Drainage lines show evidence of the action of periodically flowing water (for example, gravel, pebble, rock or sand bed, scour hole or nick point and/or incised channel at least 30cm deep).

DO NOT handle, mix, apply or conduct testing operations in areas susceptible to runoff where drainage results in rapid entry into waterways. These areas include roads, access tracks, snig tracks and compacted log dumps where no specific action has been taken to prevent runoff into waterways, or areas mounded perpendicular to the contour roads, access tracks, snig tracks and compacted log dumps.

DO NOT use or apply this product post-emergence on raised beds or where furrows have been created in the soil for the purposes of holding or channeling water.

TT Canola: DO NOT use or apply this product post-emergence on raised beds or where furrows have been created in the soil for the purposes of holding or channeling water.

SOIL MOISTURE

Imtrade Atrazine 900 WG Herbicide requires rainfall or irrigation to move it down through the soil into the weed root zone to make it effective. Sufficient rain or irrigation to thoroughly wet the soil through the weed root zone should occur or be made within 10 days after spraying.

DIRECTIONS FOR USE

Situation & Crop	Weeds Controlled	State	Rate	Critical Comments
Canola – (Triazine Tolerant (TT) varieties only) Pre-emergence or post sowing pre-emergence only	Capeweed, Charlock, Clover, Corn Cromwell, Doublegee, Fumitories, Geraniums, Ivy-leaf Speedwell, London Rocket, Mustards, Turnips, Paterson's Curse, Shepherd's Purse, Silver Grass <i>(Vulpia)</i> Suppression of Annual Ryegrass, Barley Grass, Brome Grass, Wild Oats and Wild Radish	All States	1.1 – 2.2kg/ha	Important: This use is subject to an INTEGRATED WEED MANAGEMENT STRATEGY for the use of triazine herbicides in triazine tolerant canola. See General Instructions: Integrated Weed Management Strategy for TT-Canola. Can be applied up to a week before sowing or post-sowing pre-emergence (ideally incorporated by harrows). For best results apply to bare moist soil either immediately before seeding or as a pre-emergence treatment at or within 7 days of planting. Sufficient rainfall (20-30mm) to wet the soil through the weed root zone is necessary within 2-3 weeks of application. Application should not be made to ridged or excessively cloddy soil. When applied before seeding, incorporate to a depth of 5cm.
Canola – (Triazine Tolerant (TT) varieties only) Post-emergence application	Annual Ryegrass (1-2 leaf stage only), Mustards, Wild Radish, Turnips		0.5 – 1.1kg/ha	Important: This use is subject to an INTEGRATED WEED MANAGEMENT STRATEGY for the use of triazine herbicides in triazine tolerant canola. See General Instructions: Integrated Weed Management Strategy for TT-Canola. Apply to moist soil when weeds are actively growing. The addition of 0.5-1% v/v of crop oil will enhance post-emergence activity. Cold Water: Under cold water conditions (10°C or less), use Hasten as a crop oil. DO NOT apply more than one post-emergence application.

Situation & Crop	Weeds Controlled	State	Rate	Critical Comments
Sorghum, Broom Millet, Saccaline and Forage Sorghum - DRYLAND	Amaranth, Annual Ground Cherry, Barnyard Grass, Blackberry, Nightshade, Bladder Ketmia, Burrs, Caltrop, Common Thornapple, Crowsfoot Grass, Dwarf Marigold, Fat-hen, Love Grass, Mintweed, Parthenium Weed, Pigeon Grass,	All States	2kg/ha	Pre-plant or pre-emergence only: Use this technique where grasses are likely to be the major problem.
			2kg/ha followed by 1.3kg/ha	Pre-plant or Pre-emergence followed by a Post emergence application: Use this technique where long term weed control is required because of heavy rainfall or prolonged wet conditions following the initial application; or because dry weather follows sowing and spraying and weed and grass growth occurs.
			2 – 2.5kg/ha	Post-emergence only: Use the lower rate where only a broadleaf weed problem occurs and the higher rate where grasses are the major problem. Add a non-ionic surfactant.
Sorghum, Broom Millet, Saccaline and Forage Sorghum – IRRIGATED	Pigweed, Plains Grass, Potato Weed, Prickly Paddy Melon, Spring Grass, Summer Grass, Wireweed		1.7kg/ha followed by 1.4 – 1.6kg/ha	Pre-plant or Pre-emergence followed by a Post-emergence application (See General Instructions): Use the lower rate on medium textured soils and the higher rate where grasses are the major problem or on heavy soils.
			2.5 – 3.3kg/ha	Pre or Post-emergence only (See General Instructions): Use the lower rate on light sandy soils i.e. soils low in organic matter, and medium textured soils; and the higher rate where grasses are the major problem or on heavy soils. DO NOT apply as a pre-emergence application to light sandy soils. Add a non-ionic surfactant to post-emergence applications. Note: Not recommended in the M.I.A.
	Black Bindweed (Climbing Buckwheat), Cobbler's Peg, Sesbania, Wild Oats	NSW, Vic, SA, WA only		

Situation & Crop	Weeds Controlled	State	Rate	Critical Comments
Sorghum, Broom Millet, Saccaline and Forage Sorghum – IRRIGATED & DRYLAND	Black Pigweed, Mintweed	Qld only	1.3kg/ha	Pre-plant, Pre-emergence or Post-emergence application: Apply when weeds are young (4-6 true leaves) and actively growing.
Sorghum	Parthenium Weed	All States	3.3kg/ha	Apply as a pre-emergent blanket spray. Add 150 mL of a non-ionic surfactant to each 100L spray.
	Black Pigweed, Sesbania Pea		1.1kg/ha & 400 mL/ha Amicide® 625	Post-emergence
Maize & Sweet Corn – IRRIGATED & DRYLAND	Amaranth, Annual Ground Cherry, Barnyard Grass, Blackberry Nightshade, Bladder Ketmia, Burrs, Caltrop, Common Thornapple, Crowsfoot Grass, Dwarf Marigold, Fat-hen, Love Grass, Mintweed, Parthenium Weed, Pigeon Grass, Pigweed, Plains Grass, Potato Weed, Prickly Paddy Melon, Spring Grass, Summer Grass, Wireweed		2.5 – 3.3kg/ha	Pre-plant or Pre-emergence followed by a Post-emergence application: Use the lower rate where broadleaf weeds are the major problem and the higher rate where grasses are the major problem or on heavy soils. Post-emergence application: Add a non-ionic surfactant.

Situation & Crop	Weeds Controlled	State	Rate	Critical Comments
Maize & Sorghum	Black Pigweed	Central Qld only	1.2kg/ha 350g or 1.2kg/ha	Pre-emergence. Post-emergence: Use lower rate for seedlings (2 true leaves). For plants up to 3cm in diameter use higher rate. Add a non-ionic surfactant (See Applications).
	Thornapple (<i>Datura</i> spp.) & other Broadleaf Weeds including Annual Ground Cherry, Bladder Ketmia, Caltrop, Bellvine, Mintweed, Noogoora Burr, Wild Gooseberry, Pigweed, Black Pigweed, <i>Amarantus</i> spp., Sesbania Pea, Wandering Jew	Qld, NSW only	830g – 1.1kg/ha plus 300 – 500 mL/ha Tordon 75-D	Use the lower rate when weeds are small and actively growing. Use the higher rate for larger weeds or when plants are not growing actively. Spray when the crop has 4 – 6 fully expanded leaves and secondary roots have developed. DO NOT add surfactants or crop oil. If rotating to Winter cereal crops DO NOT apply this tank mixture within 6 months of the expected sowing date. For other crops or situations observe plant back information on the respective product labels.
Lucerne	Mintweed	All States	600g/ha	Apply when the first germination of mintweed seedlings are 5-8cm high and actively growing. Using the low volume boom spray apply 140-170 litres spray mixture per hectare. DO NOT TREAT LUCERNE STANDS UNDER ONE YEAR OLD.
Oil Tea Tree	Purple Top (<i>Verbena bonariensis</i>)	All States	Post Harvest Spray 2.2kg/ha or In Crop Shielded Spray 1.8kg/ha	DO NOT apply to weeds growing in light sandy soil. Apply once per season. Add wetting agent or surfactant at label rates. Post Harvest: Apply by boom in a spray volume of at least 100L/ha immediately after harvest when the surface of stumps have dried but before bud swell. For suppression of existing plants apply in combination with Spray-Seed. In Crop: Apply using a shielded sprayer in combinations with Spray-seed or glyphosate. Avoid contact with foliage.

Situation & Crop	Weeds Controlled	State	Rate	Critical Comments
Sugar Cane	Giant Sensitive Plant	Qld, WA only	2.2 – 3.3kg/ha	<p>Apply when soil is moist. Avoid spraying weeds under stress. Apply by ground rig only and obtain good cover of target weeds. Use high rates towards the higher end of the range where heavy soil or high trash levels are present. Use higher rates for grasses.</p> <p>Approximate period of weed control: 2.2kg – up to 4 weeks. 3.3kg – up to 10 weeks. A pre-emergent application is preferred for grasses.</p> <p>Control of emerging broadleaf weeds: Where plants are large or conditions dry add 800 mL of Imtrade Smash 625 plus a non-ionic surfactant. Good coverage is essential. Use higher rate when up to 14 weeks residual control of broadleaf weeds is required.</p> <p>If grasses are a problem, then Imtrade Atrazine 900 WG Herbicide should be applied prior to weed emergence. If emerged grasses are present add Paraquat 250 Herbicide at 1.2 to 1.7L/ha.</p>
	Barnyard Grasses, Blue Top, Borreria (Square weed), Budda Pea, Cobbler's-pegs, Crowsfoot Grass, Flannel Weed, Paddy's Lucerne (Common sida), Pigweed, Sesbania, Sida retusa, Spiny Spider Flower (Wild Rose), Stinking Passion Vine, Summer Grasses, Sweet Briar, Vines	Qld, WA and NSW only		
Lupins – Weedfree Seedbed	Capeweed, Turnip, Wild Radish, Doublegee, Clovers and Medics, Mustard, Wireweed, Suppression of Annual Grass Weeds	WA only	280 - 560g/ha plus 280 - 560g/ha Simazine 900 WG	<p>Apply to bare moist soil immediately before or at seeding. Application should not be made to ridged or excessively cloddy soil. Incorporation by the sowing operation should not exceed 5cm. Sufficient rainfall (20-30mm) to wet the soil through the weed root zone is necessary within 2-3 weeks of application. Results can be variable if seasonal conditions are dry prior to sowing and lupins are sown into a dry or low moisture seedbed. Apply a maximum of 1.2kg/ha of the mixture on yellow sands.</p> <p>N.B. – Some early crop phytotoxicity may be observed particularly on yellow sands carrying native pear and pine vegetation.</p> <p>CAUTION: DO NOT use on white or grey sands.</p>

Situation & Crop	Weeds Controlled	State	Rate	Critical Comments
Eucalyptus and <i>Pinus radiata</i> plantations	Common Sowthistle, Mouse-eared Chickweed, Sand Brome, Silver Grass, Wild Oats, Yorkshire Fog Grass	All States	5 – 8.8kg/ha	<p>Pre-planting: Graze the area heavily and apply the tank mixture by boom spray or aircraft no more than two weeks before planting.</p> <p>Post-planting: Apply the mixture by boom spray either immediately after or within one month of planting. When applying by aircraft, the pines must be at least one month old. Application must be made before the Spring flush commences. Use rates towards the lower end of the range where broadleaf weeds are the major problem and rates towards the higher end of the range where grasses are the major problem. Use up to 8.8kg/ha on clay loams and heavier textured soils. For sandy soils and soils described as highly erodible, apply a maximum of 5kg/ha. DO NOT apply with a knapsack sprayer.</p>
<i>Pinus radiata</i> plantations	Wild Oats, Silver Grass, Rye Grass, Yorkshire Fog Grass, Sand Brome, Mouse-eared Chickweed, Capeweed, Clovers, Sowthistle Seedlings	ACT, Vic, SA, WA, Tas, NSW only	1.6 – 2.2kg/ha plus 5.6 – 8L ChemAg Amitrole T	<p>Pre-planting or Post-planting: Spray to pine seedlings. Use the higher rate where grasses are a major problem. Preferably apply as a strip, rather than a blanket application. Method of application: Refer Application Table.</p>

Situation & Crop	Weeds Controlled	State	Rate	Critical Comments
Grass Seed Crops (Established stands of Sirocco, Phalaris, Demeter Fescue & Currie Cocksfoot), Seedling Signal Grass & <i>Panicum maximum</i>	Brome Grass	All States	1.0 – 1.2kg/ha	Apply after the Autumn break when Brome Grass is just emerging. Apply by low volume boom spray.
	Billygoat Weed (Blue Top), Cobbler's-pegs, Crowsfoot Grass, Fleabanes, Love Grass, Mexican Poppy, Setaria spp., Sida spp., Stinking Roger, Thickhead, Wild Hops, Wild Radish, Woolly Top, Rhodes Grass	Qld only	2.5 – 3.3kg/ha	Pre-emergence: Apply at or immediately after planting, preferably to a moist soil and before crop and weeds germinate. Use the lower rate on <i>Panicum maximum</i> seed crops and the higher rate on Signal Grass when grasses are likely to be the major problem. Guinea Grass, Panics and Green Summer Grasses (<i>Brachiaria</i> spp.) are not controlled. WARNING: Damage can occur to <i>Panicum maximum</i> when application is made under cool, dry conditions. NOTE: When used on Signal Grass and <i>Panicum maximum</i> , user accepts all responsibility should any damage occur.
Seedling Ryegrass, Seed Crops	Winter Grass, Toad Rush	Vic only	480 – 550g/ha	Use the lower rate at 2-3 leaves and the higher rate at early tillering.
	Winter Grass, Toad Rush, Broadleaf Weeds		550g/ha plus 500 mL/ha MCPA 500 & 160 mL/ha Kamba® 500	Use at the tillering stage.
Established Ryegrass, Seed Crops	Winter Grass, Toad Rush, Loosestrife, Sorrel, Barley Grass, Docks, Soft Brome, Silver Grass		830g – 1.1kg/ha	Apply after good Autumn rains have fallen and conditions are becoming cooler - around late May to mid-June - graze heavily prior to application NOTE: Some crop damage may occur on sandy soils or if crop is water stressed. DO NOT apply Nitrogen prior to spraying.
Fallow Area Maintenance (Prior to sowing Wheat, Peas, Lupins)	Broadleaf Weeds and Grasses	Vic only	650g – 870g/ha	Apply late July/mid-September for May-June sowing. Use the higher rate for a 14 month fallow – apply in Feb/April before Autumn rains.

Situation & Crop	Weeds Controlled	State	Rate	Critical Comments
Fallow Area Maintenance prior to planting a Sorghum crop in a conservation tillage system	Amaranth, Annual Ground Cherry, Barnyard Grass, Black Bindweed (Climbing Buckwheat), Blackberry, Nightshade, Bladder Ketmia, Burrs, Caltrop, Common Thornapple, Crowsfoot Grass, Dwarf Marigold, Fat-hen, Love Grass, Mintweed, Parthenium Weed, Pigeon Grass, Plains Grass, Potato Weed, Prickly Paddy Melon, Sesbania Pea, Spring Grass, Summer Grass, Wireweed	ACT, Qld, NSW only	2 – 3.3kg/ha	Pre-plant: Apply to moist soil or when rain is imminent and prior to the germination of weeds and grasses, or as a tank mixture with specific knockdown herbicide if weeds and grasses are present. Use the lower rate when short-term control of weeds and grasses is required or where a wheat crop will follow sorghum in the rotation. Use the higher rate when longer term control of weeds and grasses is required.
Roadside and Rights of Way	Parthenium Weed	ACT, Qld, NSW & NT only	3.3kg/ha	Pre-emergence and Post-emergence: Apply to moist soil following germinating rains and when further follow up rain is imminent. Where germination has occurred, ensure application is made to seedling plants.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

HARVEST:

ALL CROPS: NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING:

CANOLA: PRE-EMERGENCE APPLICATION: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 15 WEEKS AFTER APPLICATION.

CANOLA: POST-EMERGENCE: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 6 WEEKS AFTER APPLICATION.

OTHER CROPS (EXCEPT CANOLA): DO NOT APPLY TO AREAS THAT WILL OR MAY BE GRAZED OR CUT FOR STOCK FOOD WITHIN 28 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions.

Imtrade Atrazine 900 WG Herbicide is a pre- and post-emergence herbicide, which will selectively control:

- Weeds and grasses in Canola (Triazine Tolerant (TT) varieties only), Sorghum, Maize, Sweet Corn, Sugar Cane, Lupins, Broom Millet, Saccaline and Forage Sorghum.
- Mintweed in established Lucerne.
- Brome Grass in Sirocco Phalaris, Demeter Fescue and Currie Cocksfoot grass seed crops.
- Provide control of weeds and grasses growing on a fallow in a conservation tillage system.

However, established perennial species, large annuals and large broadleaf weeds are not satisfactorily controlled at the rates recommended. It acts mainly through root absorption, its effectiveness depends on the occurrence of rainfall or irrigation to move it down onto the weed root zone. Duration and effectiveness of control depends on the amount of chemical applied, soil type, rainfall and particular weed species.

The maximum rate of atrazine application in all crops except plantation forestry is limited to an amount of product equivalent to 3kg a.i. atrazine/ha per year. DO NOT exceed this limit, especially when applying an atrazine herbicide post-emergence, where an atrazine herbicide has been applied pre-emergence.

The maximum rate of application in plantation forestry is an amount of product equivalent to 4.5kg a.i. atrazine/ha per year in sandy soils and those defined as highly erodible, and product equivalent to 8kg a.i. atrazine/ha per year in clay loam and heavier textured soils.

MIXING

- Fill spray tank 60-80% full with clean water BEFORE adding Imtrade Atrazine 900 WG Herbicide.
- Begin agitating tank contents vigorously and continue agitation during entire mixing and spraying operations.
- Pour required amount of Imtrade Atrazine 900 WG Herbicide STEADILY in the spray tank. Allow vigorous bypass agitation to completely disperse product. **DO NOT** dump product into spray tank all at once.
- After adding required quantity of Imtrade Atrazine 900 WG Herbicide and obtaining COMPLETE DISPERSION, continue to fill tank to desired level for spraying.
- Thorough agitation of the spray liquid should continue during the entire spraying operation. **NB:** Spray solution should **NOT** be left standing in the tank overnight.
- **DO NOT** mix, load or apply this product within 20m of any well, sink hole, intermittent or perennial stream.

INCORPORATION (PRE-PLANT AND AT SOWING APPLICATION)

Imtrade Atrazine 900 WG Herbicide acts mainly by root absorption. Its effectiveness depends on the occurrence of rainfall or irrigation to move it down into the weed root zone. Sufficient rain or irrigation to thoroughly wet the soil through the weed root zone should occur or be made immediately after application to provide appropriate weed control. Delay in activation of Imtrade Atrazine 900 WG Herbicide may result in some weed growth.

In floor or furrow irrigation situations, complete and continued activation of the product may not occur due to a thin band of dry soil on the surface during or after irrigation. Mechanical incorporation after application, using light harrows to incorporate the product into the soil not more than 4cm deep, is required to ensure the irrigation water activates the product. Always apply Imtrade Atrazine 900 WG Herbicide to an even unridged seedbed.

Integrated Weed Management Strategy for TT-Canola

An Integrated Weed Management Strategy (IWM) for TT-Canola (the Strategy) has been developed by Imtrade Australia Pty Ltd with the assistance and agreement of the Canola Association of Australia. The Strategy outlines recommendations, measures and options of weed management including management of herbicide resistance in weed populations. The Strategy is available from an Imtrade Australia Pty Ltd representative and the Canola Association of Australia. A program has been developed that outlines sound agronomic practices and IWM programs designed to optimize the performance of TT-Canola. It is advised that consultation on IWM be undertaken with an accredited agronomist prior to the use of Imtrade Atrazine 900 WG Herbicide on TT-Canola.

To minimize herbicide resistance:

- Avoid dry sowing in heavily weed-infested paddocks. Wait for a weed germination after the opening rains in weedy paddocks. Use a pre-plant knockdown or cultivation. No weeds should be allowed to survive at this stage.
- Adapt the weed control program to the anticipated weed spectrum and pressure:
- **Broadleaf Weeds and Ryegrass:** Use Simazine or Atrazine plus Imtrade Trifluralin 480 pre-emergence. A follow-up with a Group 1 Herbicide (if ryegrass is susceptible) or Atrazine may be necessary.
- **Broadleaf Weeds only:** Use ATRAZINE post-emergence.
- **DO NOT** use Atrazine or Simazine if the area to be treated has had a triazine herbicide applied to it last season.
- Watch for escapes in paddocks with a long history of Group 5 Herbicide use.
- **DO NOT** use Group 5 Herbicides in consecutive years.

Resistant Weeds Reporting

Growers should collect plant or seed samples where weeds that are normally susceptible to Atrazine and Simazine may be resistant, get them tested and seek professional advice.

To avoid Triazine carry-over

On acid soils (pH less than 6.5) - The maximum rate of Imtrade Atrazine 900 WG Herbicide or Simazine 900 WG Herbicide or a combination of the two products to be applied to the crop during the growing season is 2.2kg/ha.

On alkaline soils (pH greater than 6.5) - The maximum rate of Imtrade Atrazine 900 WG Herbicide or Simazine 900 WG Herbicide or a combination of the two products to be applied to the crop during the growing season is 1.1kg/ha.

Post-emergence use - It is recommended that Imtrade Atrazine 900 WG Herbicide only be used, and at rates of 1.1kg/ha or less, on both acid or alkaline soils.

SUMMER CROPS

Heavy rains immediately following an application tend to result in excessive concentrations of herbicide in the seed furrow, thus encouraging possible crop injury. This is most likely to occur when a pre-plant or pre-emergence application is made using rates in excess of 2kg/ha. To avoid the likelihood of herbicide injury follow time of application (a), (b) or (d) listed below:

Time of Application (Sorghum, Maize, Broom Millet, Saccaline, Sweet Corn).

a) Pre-plant Application – followed by Post-emergence Application.

Apply after establishment of the seedbed and up to two weeks prior to sowing. Application should preferably be made to moist soil. If using the technique on irrigated crops then a post-emergence spray must be applied as soon as possible after the crop has emerged and has developed 2 to 3 leaves.

b) At Sowing Application – followed by a Post-emergence Application.

Apply at or immediately after planting and before crop and weeds emerge. Application should preferably be made to moist soil. If using the technique on irrigated crops then a post-emergence spray must be applied as soon as possible after the crop has emerged and has developed 2 to 3 leaves.

c) At Sowing Application.

Apply at or immediately after planting and before crop and weeds emerge. Application should preferably be made to moist soil, and rain or irrigation should follow application. Mechanical incorporation may also assist.

d) Post-emergence Application.

Application must be made to seedling broadleaf weeds and grasses when they are not more than 1cm high. Normally the crop is then at the 2 to 3 leaf stage. For best results, the soil should be moist and rain or irrigation should follow application. Add a non-ionic surfactant for all post-emergence applications in Sorghum, Broom Millet and Saccaline, and a crop oil at the recommended rate to the spray mixture for Maize and Sweet Corn.

APPLICATION

Ground Application: Application can be made as an overall or band treatment. Minimum bandwidth should be 30cm. Apply 50 to 100 litres per hectare.

Aircraft Application: With aircraft application the need for good soil moisture at time of application and follow up rain or irrigation within 10 days is most critical. Apply 10 to 20 litres per hectare. **DO NOT** use human flaggers in aerial spraying, unless protected by engineering controls such as enclosed cabs.

TT-Canola: DO NOT apply to TT-Canola by aircraft. Apply only with a low boom sprayer with a 60m buffer zone downwind of treated fields to natural or impounded lakes or dams, and a 20m buffer zone for any well, sink hole, intermittent or perennial stream. Apply only to areas where run-off is unlikely to occur or where run-off may be captured by farm earthworks. Application can be made as an overall spray or as a band spray. Minimum bandwidth should be 30cm.

METHOD OF APPLICATION TABLE

Pinus radiata plantations

Pre-planting

Type of Application	Critical Comments
Pre-planting	Graze area heavily and apply the tank mixture no more than 2 weeks before planting.
Post-planting	Apply tank mixture either immediately after or within one month of planting. Application must be made before the Spring commences. Avoid spraying the pine seedling by using directed spray.

COMPATIBILITY

Imtrade Atrazine 900 WG Herbicide is compatible with Paraquat 250 Herbicide and other residual herbicides such as Diuron. In Sugar Cane only, addition of Sodium 2,4-D will assist in short term suppression of Nutgrass and other sedges.

RESISTANT WEEDS WARNING

GROUP	5	HERBICIDE
-------	----------	-----------

Imtrade Atrazine 900 WG Herbicide is a member of the triazines group of herbicides. Imtrade Atrazine 900 WG Herbicide has the inhibition of photosynthesis at photosystem II mode of action. For weed resistance management, Imtrade Atrazine 900 WG Herbicide is a Group 5 Herbicide. Some naturally-occurring weed biotypes resistant to Imtrade Atrazine 900 WG Herbicide and other Group 5 Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These weeds will not be controlled by Imtrade Atrazine 900 WG Herbicide or other Group 5 Herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Imtrade Australia Pty Ltd accepts no liability for any losses that may result from failure of Imtrade Atrazine 900 WG Herbicide to control resistant weeds. Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture/Primary Industries, or an Imtrade Australia Pty Ltd representative. Any incidents of resistance must be reported to Imtrade Australia Pty Ltd, Toll Free 1800 171 799.

PRECAUTION

Re-entry period: DO NOT enter treated areas without protective clothing until spray has dried.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray foliage of desirable plants.

DO NOT apply high rates of application to heavier soils if roots of desirable shrubs or trees are near the surface.

DO NOT use in channels and drains.

DO NOT use near newly planted shrubs, young ornamentals and species with shallow roots e.g. Prunus species, or trees in sandy porous soils.

DO NOT apply under weather conditions or from spraying equipment that may cause spray drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT apply or drain or flush equipment on or near areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots.

DO NOT plant crops other than those recommend on this label for at least 6 months following treatments at rates up to 1.4kg/ha and for 18 months following treatments of 1.4 to 3.3kg/ha.

DO NOT apply in excess of 3.3kg product/ha in any one year, except in forestry situations.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT apply this product within 60m of natural or impounded lakes or dams.

DO NOT contaminate dams, waterways or drains with this product or used containers.

DO NOT apply under weather conditions or from equipment that may cause drift of this product or spray mix into adjacent areas, particularly wetlands, waterbodies or watercourses.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area out of direct sunlight. Shake empty bag into spray tank. Single rinse plastic bags before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. Puncture or shred and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. **DO NOT** inhale dust or spray mist. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), a washable hat and elbow-length PVC gloves. If using a hand directed sprayer, wear in addition: waterproof trousers and boots. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (*Phone Australia 13 11 26; New Zealand 0800 764 766*).

Additional GHS Hazard & Precautionary Statements

•May cause an allergic skin reaction; •May cause damage to organs through prolonged or repeated exposure; •Very toxic to aquatic life with long lasting effects; •Avoid breathing fumes, mists, vapours or sprays; •**DO NOT** get on clothing; •Contaminated work clothing should not be allowed out of the workplace; •Avoid release to the environment; •Get medical advice or attention if you feel unwell; •**IF SWALLOWED:** Rinse mouth. Do **NOT** induce vomiting; •**IF ON SKIN:** Wash with plenty of soap and water; •If skin irritation or rash occurs: Get medical advice; •Collect spillage; •Not combustible. Use extinguishing media suited to burning materials. Water fog or fine spray is the preferred medium for large fires; •Store in a dry place.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or from Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

UN 3077	ENVIRONMENTALLY HAZARDOUS SUBSTANCES, SOLID, N.O.S. (contains ATRAZINE)
IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	
PG III	HAZCHEM 2Z



APVMA APPROVAL No: 63503/122726