


IMTRADE
JIGSAW[®]

800 WG FUNGICIDE

Get rid of Grey

- ▶ Effective protection against Grey Mould
- ▶ 1 Day withholding period
- ▶ Long-life granular formulation



IMTRADE JIGSAW® 800 WG FUNGICIDE

Jigsaw® Benefits

Effective protection - Jigsaw® offers unique target site action belonging to Group 17 Sterol Biosynthesis Inhibiting fungicide.

Short withholding period - 1 day withholding period in Strawberries.

Long-life formulation - Jigsaw® is a powerful 800 g/kg granular Fenhexamid formulation, offering extended shelf-life.

Botrytis Resistance Management - Rotate Jigsaw® regularly with other chemical groups between consecutive sprays to prevent resistance development.

IPM Friendly - Jigsaw® fits into IPM, being compatible and not affecting Botrytis antagonists such as Trichoderma and Ulocladium.

Grapevine Management

Primary infection site for grapes is at the cap-scar created by cap fall at flowering. Jigsaw® has demonstrated to be most efficacious in reducing overall orchard Botrytis infection if applied as a preventative measure at flowering.

3 spray program*	Jigsaw® once only
4 spray program*	Max. of 2 sprays of Jigsaw®

***DO NOT** apply in volumes less than 250 L/ha.

- Monitor the crop application at full bloom for optimal control
- Apply to the point of run-off
- Volumes of 500-1500 L/ha (canopy size dependent)
- Good coverage of bunches is key
- WHP: 21 days (domestic market); Export wine grapes: Consult your winery, industry spray diary, or peak industry body for recommended withholding period. Do not apply later than 80% capfall.

Grey Mould Management

Strawberry infection happens primarily during flowering stage as the pathogen enters soft tissues of the flowers where it either colonizes infected flowers or enters a state of dormancy. Botrytis then infects the fruit from the receptacle of the developing Strawberry or where petals are trapped against the fruit. Lesions could then develop from infection even until post-harvest and if left uncontrolled could infect neighbouring fruits.

If conditions favour disease development:

- 7 to 10 day interval between sprays
- **DO NOT** apply more than 2 successive sprays of Jigsaw®
- **DO NOT** apply volumes less than 500 L/ha



For more information on Jigsaw®, visit [imtrade.com.au](https://www.imtrade.com.au)
or contact your local Imtrade representative on **1800 171 799**

ALWAYS READ AND FOLLOW LABEL DIRECTIONS BEFORE USING ANY PRODUCT IN THIS FLYER.
COPYRIGHT Information contained herein remains the copyright of Imtrade CropScience
and cannot be reproduced without the company's permission. © 2023